



Northern New York Agricultural Development Program

FACT SHEET

5 in a series on Phosphorus

Achieving Yield and Quality with Less P

**Are band-
applications of
fertilizer P needed
to achieve
optimum yield
and quality of
corn for silage on
fields that test
high or very high
in P?**

**Researchers
working with
Northern New
York farms have
answers for this
question.**

The Impact of Starter P on Corn Silage Quality

Principal Investigators: Dr. Quirine Ketterings, Assistant Professor of Crop & Soil Sciences, Cornell University; Karl Czymmek, Senior Extension Associate, Pro-Dairy Program, Cornell University

Introduction:

Optimizing Silage Yield and Quality

Due to increasing pressure on dairy producers to reduce the amount of phosphorus (P) applied to soils in the form of fertilizer, Cornell researchers conducted on-farm and research station trials from 2002 through 2004 to determine if band-applications of fertilizer P are needed for optimum yield and quality of corn for silage on fields that test high or very high in P.

Excess phosphorus in field runoff into nearby surface waters can cause eutrophication, an algae bloom that disturbs the aquatic habitat. If farmers can reduce the amount of P applied to fields without losing crop yield or quality, the risk of this environment disturbance decreases.

Methods

From 2000 to 2004, sixteen trials set on Northern New York farms were analyzed. Trials were set on fields testing high or very high (more than 8 lbs P/acre Morgan extractable P/acre) for P. Plots were harvested for corn silage.

Soil samples were analyzed for each plot at planting and at PSNT time when corn reached 6 to 12 inches tall. Application for the on-farm trials included 1) no starter applied, 2) no P in the starter (N+K only), 3) low or recommended P in starter (10-25 lbs of P₂O₅ per acre) and 4) producer's starter blend and rate.

The four methods evaluated for research station trials were:

- no starter applied
- no P in the starter (200 lbs of 10-0-10)
- 20 lbs of P₂O₅ in the starter (200 lbs of 10-10-10)
- 40 lbs of P₂O₅ in the starter (200 lbs of 10-20-10).

**See Fact Sheet #4:
Limiting Phosphorus
Use for Corn
Growing in NNY for
more details on when
less P is better for
growing corn**

**Want to know
more about
phosphorus use?**

The following Fact Sheets are available from the Northern New York Agricultural Development Program and your local Cornell Cooperative Extension office:

- Why is Phosphorus an Issue for New York State Farms?
- Trends in Soil P Status
- Developing a P Index for NNY Soils
- Limiting Phosphorus Use for Corn Growing in NNY Without Sacrificing Yield
- The Impact of Starter P on Corn Silage
- P Runoff: Calibrating the P Index for NNY

All research station trials were done in four replicates. Most on-farm trials were conducted in two replicates. Replications, or repeated measurements in the same field, are important to improve confidence in test results.

**Results:
Impact on Silage Yield**

Research results indicate that applying starter P on sites that test very high and when manure is applied to sites that test high is unlikely to result in a yield response, so starter P can be safely eliminated on these sites. On sites that test high and have no manure applications P starter levels can be reduced to less than 25 lbs P₂O₅/acre without affecting yield. See Table 1 for silage yields (tons/acre 35% dry matter) for on-farm trials conducted in 2001-2003.

Table 1. Silage yields (tons/acre 35% dry matter) for on-farm trials conducted in 2001-2003.

	# of trials	2001	2002	2003	3-Year Average	
		(27)	(16)	(22)	High P	Very High P
No starter		16.7 b	15.7 a	20.6 b	17.7 b	19.5 b
N(+K) only		19.3 a	16.2 a	20.7 b	17.9 b	20.6 ab
N(+K) + 10-25 lbs P ₂ O ₅ /acre		19.9 a	16.5 a	21.7 a	19.2 a	21.4 a
N(+K) + > 25 lbs P ₂ O ₅ /acre		19.8 a	16.0 a	21.1 ab	18.2 ab	21.2 a

Key: a, b = average values within columns are statistically different (α=0.05)

Impact on Silage Quality

Table 2 shows the results of the quality analyses for the study. Bottom line is that differences were not significant and were well within laboratory analytical uncertainty, indicating that leaving P out of the starter fertilizer in high or very high P soils did not impact silage quality.

Conclusions

With increased attention directed toward P non-point source pollution, it makes little sense to use more starter P than is necessary to support optimum yields, especially on fields where significant amounts of manure nutrients are regularly applied. Corn responds to nitrogen (N) in the starter band more often than P and we continue to recommend 20-30 lbs of banded starter N, even where P is eliminated.

We suggest corn growers test fields for soil fertility status at least once in three years, apply manure to low and medium P fields, and adjust starter P application rates accordingly (Table 3).

Table 2. Effect of starter P addition on corn silage quality in New York State.

Quality parameter	Research Station Trials (9)				On-Farm Trials (62)			
	No Starter	200 lbs 10-0-10*	200 lbs 10-10-10*	200 lbs 10-20-10*	No Starter	N(+K) only	N(+K)+ 10-25 lbs P ₂ O ₅ /acre	N(+K)+ > 25 lbs P ₂ O ₅ /acre
	----- % of dry matter -----							
Moisture content	64	65	64	64	60	60	60	59
NDF+	43.5	42.3	43.4	42.8	42.1	42.6	42.7	41.6
	----- % of NDF -----							
Digestibility of NDF	62.2	62.2	62.2	62.6	62.3	60.8	61.7	61.6
	----- lbs -----							
Milk/ton of silage	3692	3699	3700	3703	3734	3652	3683	3712
	----- % of dry matter -----							
Crude protein	7.4	7.3	7.3	7.5	7.6	7.5	7.7	7.6
P	0.21	0.21	0.20	0.20	0.23	0.23	0.23	0.23
K	0.83	0.88	0.85	0.84	1.09	1.09	1.10	1.11
Ca	0.21	0.20	0.19	0.20	0.17	0.18	0.18	0.18
Mg	0.19	0.17	0.17	0.18	0.14	0.14	0.14	0.1
	----- % of dry matter -----							
Zn	16.4	16.6	16.0	15.6	17.6	17.9	17.3	16.5
Cu	4.3	4.3	4.2	4.3	3.8	3.9	3.7	4.1
Mn	17.6	16.9	16.2	16.8	13.3	13.7	13.6	13.7

*Key: * per acre; + NDF: Neutral Detergent Fiber (48hr)*

Table 3. Phosphorus fertilizer guidelines for corn in New York State.

Soil Test P	lbs P ₂ O ₅ /acre	
	With manure	No manure
Very Low	20-30	60-70*
Low	20-30	50-60*
Medium	20-30	25-50*
High	0	0-25
Very High	0	0

*Key: * Put ~25 lbs P₂ O₅/acre in the starter fertilizer band; balance may be included in the band or broadcast.*

New York Starter P Project Sponsors

The New York State Starter Phosphorus Project was funded by a research and extension grant from the Northeast Sustainable Agriculture and Education Program and contributors from the Natural Resource Conservation Service and the Northern New York Agricultural Development Program Program. Fertilizer for the on-farm and station trials was donated by Agway's Lyon blend plant and Carovail; seed was donated by Pioneer Hi-Bred International, Inc.

Principal Investigators

Dr. Quirine Ketterings, Assistant Professor, Nutrient Management Spear Program, Department of Crop and Soil Sciences, Cornell University; and Karl Czymmek, Senior Extension Associate, ProDairy, Department of Animal Science, Cornell University

Project Coordinator: Sheryl Swink, Program Support, Nutrient Management Spear Program, Department of Crop and Soil Sciences, Cornell University

Project Assistants: Greg Godwin, Resource Support Specialist; Greg Albrecht, Extension Associate, Cornell University

Participating Farmers

Canton: Jon Greenwood, Kevin McCollum; Carthage: John Williams; Chase Mills/Waddington: Gary Tiernan; Lisbon: Ken Pemberton; Madrid: David Fisher; Paul, Tim and Mark Heiden; and Lou Ann King.

Participating CCE Educators and Research/Education Institute Representatives:

Cornell Cooperative Extension of Jefferson County: Mike Hunter, 315-788-8450; Cornell Cooperative Extension of Lewis County: Jen Beckman, 315-376-5270; Cornell Cooperative Extension of St. Lawrence County: Peter Barney, 315-379-9192; William H. Miner Institute, Chazy: Everett Thomas, 518-846-7121; Willsboro Agricultural Research Farm: Mike Davis.

For more information on the Starter P project, contact: your local Cornell Cooperative Extension office; Dr. Quirine Ketterings, Nutrient Management Spear Program, Cornell University, qmk2@cornell.edu, 607-255-3061, <http://nmsp.css.cornell.edu/projects/starterp.asp>; or Karl Czymmek, kjc12@cornell.edu, 607-255-4890.

The Northern New York Agricultural Development Program selects and prioritizes research the results of which can be practically applied to farms in the six-county region of Northern NY: Jefferson, Lewis, St. Lawrence, Franklin, Clinton and Essex Counties.

To learn more about the Northern New York Agricultural Development Program, contact Co-Chairs Jon Greenwood, 315-386-3231, or Joe Giroux, 518-563-7523; or R. David Smith, Cornell University, 607-255-7286; or visit www.nnyagdev.org. ♦

4: P and Corn Silage



Northern New York
Agricultural Development Program

Northern New York
Agricultural Development
Program
162 Morrison Hall
Cornell University
Ithaca, NY 14853
607-255-7286
www.nnyagdev.org