

Northern New York Agricultural Development Program 2009 Project Report

Improved Apple Orchard Management Systems and Rootstocks for Northern NY

Project Leader

Terence Robinson, Dept. of Hort. Sciences, Cornell University, Geneva, NY 14456, ph: 315-787-2227, e-mail: tlr1@cornell.edu

Collaborator

Kevin Iungerman, Cornell Coop. Ext., 50 West High Street, Ballston Spa, NY 12020, ph 518-885-8995, email: kai3@cornell.edu

Grower Cooperators

Tom Everett, Everett Orchards, Peru NY
Donald Green III, Chazy Orchards, Chazy, NY
Seth Forrence, Forrence Orchards, Peru, NY
Mac Forrence, Forrence Orchards, Peru, NY
Adam Sullivan, Sullivan Orchards, Peru, NY
Hugh Gunison, Gunison Orchards, Crown Point, NY

Introduction

The Northern New York (NNY) apple industry is large (4,000 acres and a farm gate value of \$16 million) and is an important segment of Northern New York agriculture.

The industry has knowledgeable and progressive growers, an extensive infrastructure, and proximity to large markets. However, to remain competitive in the world apple market NNY apple growers need to continue to modernize their orchards to improve orchard production efficiency and fruit quality.

Modern high-density orchard planting systems, will help improve efficiency, yield and fruit quality and will offer growers the opportunity to plant profitable new varieties. Replanting older orchards to new high-density orchards with popular new varieties will help the long-term viability of the Northern New York apple industry.

The goal of this project was to develop and extend to growers information on modern, competitive orchard systems that incorporate new high priced varieties, disease resistant rootstocks, high planting densities for early production and partial labor mechanization to reduce costs.

Research results on high density orchards and new rootstocks conducted in other parts of NY state is not directly transferable to the colder climate of NNY.

We have utilized on-farm orchard systems and rootstock experiments that the project leaders have already established in NNY. In addition new on-farm experiments were conducted in 2009 on improved chemical thinning with Honeycrisp, drop control strategies McIntosh and fruit quality with McIntosh and Honeycrisp.

The project involved all of the apple growers in NNY through field days, workshops and winter fruit grower meetings.

Materials and Methods

We had previously established 4 on-farm trials in Clinton County that were used in this research project.

- 1) Chazy Orchards 2001 Rootstock Trial. This replicated field plot compares 16 rootstocks (G.16, G.30, B.9, B.118, O.3, Vineland 1, Vineland 3, Supporter 4, Mark, M.9T337, M.9Nic29, M.9/MM.111, M.26, M.7, MM.106, and MM.111) for survival, productivity, and adaptability to the cold climate of NNY with Honeycrisp and McIntosh as the scion varieties. The experimental design is a randomized complete block 8 replications and 10 trees per experimental unit. We measured yield, fruit size and survival for each of the rootstocks. We will publish the final report on this trial at the end of 2010.
2. Everett Orchards 2002 Orchard Systems Trial. This replicated field plot was established at Everett Fruit Farm in Peru, NY and it compares 5 orchard system (Central Leader on MM.111, Slender Pyramid on M.26 and G.30, Vertical Axis on M.9, B.9 and G.16, Solaxe on M.9, B.9 and G.16 and Tall Spindle on M.9, B.9 and G.16). The objective of the trial was to develop realistic performance and cost data for the colder part of NY state to provide growers with practical examples of different orchard system performance and economics. Densities range from 218 trees/acre to 1307 trees/acre. Varieties include McIntosh and Honeycrisp. The experimental design is a randomized complete block split plot with 3 replications and 30 trees per experimental unit. We measured yield, fruit quality, and labor input requirements for each of the various tree forms and planting densities. We will perform an economic analyses of the trial utilizing the actual packout and labor costs in 2 more years when the trial is 10 years old.
- 3) Forrence Orchards 2002 CG Rootstock Trial: This replicated field plot compares 17 new rootstocks from the Geneva apple rootstock breeding program and 8 Malling stocks from England, 2 stocks from Russia, Ott.3 from Canada, P.22 from Poland and Supporter 4 from Germany with Honeycrisp as the scion. This trial is a comparison of many of the new disease resistant rootstocks from Cornell which have substantial potential in NNY. The experimental design is a randomized complete block 10 replications and 1 tree per experimental unit. We measured yield, fruit size and survival for each of the rootstocks.
- 4) Forrence Orchards 2008 CG Rootstock Trial: A new replicated field was planted with Mac Forrence at his Valcour farm which compares 34 new rootstocks from the Geneva apple rootstock breeding program and 3 Malling stocks from England, B.9 from Russia, Ott.3 from Canada, P.22 from Poland and Vineland 1 from Canada with Honeycrisp as the scion. This trial is a comparison of many of the newest disease resistant rootstocks from Cornell which have substantial potential in NNY We measured tree survival for each of the rootstocks in 2009. It will have its first crop in 2010.

In addition, we established 3 one-year thinning, return bloom management trials with Honeycrisp and a pre-harvest drop control trials with McIntosh apple in 2008.

- 1) Thinning and Return Bloom of Honeycrisp (Chazy): In 2009 we conducted a replicated field study at Chazy orchards of timing and concentration of chemical thinners to managed cropload and return bloom on the new highly priced apple variety, Honeycrisp. This variety is proving to be difficult to manage and improved thinning strategies are essential to the long-term success of this variety. This study evaluated single vs. multiple sprays of NAA/Sevin for thinning efficacy and summer NAA sprays for improved return bloom of Honeycrisp. The experimental design was a randomized complete block with 4 replications and 2 trees per experimental unit.

2) Control of pre-harvest drop with McIntosh (Chazy): We conducted a replicated field trial where we evaluated ReTain, and NAA in 2009 to reduce pre-harvest drop of McIntosh. The trial was conducted at Chazy Orchards in cooperation with Tre Green. The objective was to determine the effect of Retain, or Retain combined with NAA, on preharvest drop of McIntosh apples in the Champlain Valley. The treatments were:

Trt No.	Treatment	Date of Application
1.	Untreated Control	
2.	Retain 333 g/acre plus Silwet L-77 0.1% v/v	28 days before anticipated harvest
3.	NAA 20 ppm	14 days before anticipated harvest
4.	Retain 333 g/acre plus Silwet L-77 0.1% v/v plus NAA 20 ppm	28 days before anticipated harvest
5.	Retain 333 g/acre plus Silwet L-77 0.1% v/v NAA 20 ppm	28 days before anticipated harvest 14 days before anticipated harvest
6.	Retain 333 g/acre plus Silwet L-77 0.1% v/v plus NAA 20 ppm	14 days before anticipated harvest
7.	Retain 167 g/acre plus Silwet L-77 0.1% v/v plus NAA 20 ppm	14 days before anticipated harvest

Results and Discussion

Orchard Systems Study (Table 1, Figures 1-8):

Our comparison of 5 orchard production systems has shown that the high density Tall Spindle system has been the most productive system in the Champlain Valley.

The Tall Spindle had the earliest production with a small crop in the second year (Figs 1 and 2). The M.9 trees had more yield than either B.9 or G.16. M.26, G.30 and MM.111 had no crop in the second year. In the third and fourth years there was a linear relationship of density and yield with the M.9 rootstock having greater yield than any of the other stocks (Fig 5 and Fig. 6). In the fifth year (2006) frost and poor pollination reduced crop significantly with McIntosh but not with Honeycrisp. However, Honeycrisp suffered from biennial bearing and had less than a full crop. B.9 rootstock was the most productive rootstock with Honeycrisp in 2006 but M.9 and G.16 were the most productive with McIntosh. In 2007 and 2008 there was a large crop with both varieties. In 2009 there was the largest crop top date with both McIntosh and Honeycrisp. The tall spindle system again had the highest yield and with McIntosh on either M.9 or B.9 rootstocks, yields reached 1600 bushels/acre (Table 1). With Honeycrisp the most productive combination was the Tall Spindle on either M.9 or B.9 rootstock which had a yield of 1200 bushels/acre. There was little difference in yield between the Vertical Axis and the SolAxe systems. The lowest yielding system was the Central Leader.

The differences in yield between systems was largely the result of the planting density. At the end of 8 years, there was a strong positive linear relationship between tree planting density on cumulative yield (Fig.3). The Central Leader system which had the lowest tree density had the lowest yield, followed by the Slender pyramid, Vertical Axis, SolAxe and Tall Spindle.

Among rootstocks M.9 has the highest yield in most years with McIntosh followed by B.9, G.16, G.30, M.26 and MM.111 (Fig 7). With Honeycrisp, B.9 had the greatest cumulative yield followed by M.9, G.16, G.30, M.26 and MM.111 (Fig. 7). B.9 and G.30 rootstocks had less biennial bearing than either M.9 or G16.

Crop value was greatest with the tall spindle system in each year except 2006 when frost damage reduced crop value with the Tall Spindle more than any other system. Nevertheless, in 2007, 2008 and 2009 the tall spindle again had the greatest crop value. The Tall Spindle had the greatest cumulative crop value followed by the Vertical Axis and SolAxe which did not differ significantly, then the Slender Pyramid and lastly the Central Leader (Fig 4 and Fig. 8). The Tall Spindle exceeded the cumulative crop value of the Central Leader by 7.7 fold with McIntosh and 10 fold with Honeycrisp.

Honeycrisp trees on all rootstocks and systems yielded less than McIntosh (80% yield) but the cumulative crop value of Honeycrisp yield was 4 times that of McIntosh due to higher fruit price and larger fruit size. By the end of the 8th year the best Honeycrisp system had accumulated \$84,347 in cumulative crop value compared to only \$19,396 for McIntosh. This level of returns would essentially pay for the establishment cost of the Honeycrisp block by the end of the 5th year but it is likely to take 10 years with McIntosh.

This trial shows that much higher yields than previously thought possible can be achieved with the Tall Spindle system at a relatively young orchard age. This high yielding system when coupled with a high priced variety like Honeycrisp can dramatically change the orchard profitability potential for new orchards in NNY State.

Table 1. Performance of McIntosh and Honeycrisp apple trees on 6 rootstocks trained to 5 orchard systems in the Champlain Valley.

Variety	System	Stock	Yield/ Acre 2009 (bu)	Crop Value/ 2009 (\$/ acre)	Cum Yield/ Acre (bu)	Av Fruit Size (g)	Cum Crop Value (\$/ acre)
Honeycrisp	Central Leader	MM.111	326	8375	546	231	12,939 e*
	Slender Pyramid	G.30	1089	25924	2472	236	55,541 bc
	Slender Pyramid	M.26	591	14146	1530	217	33,771 d
	SolAxe	B.9	817	20494	2449	230	56,617 bc
	SolAxe	G.16	796	19950	2382	231	54,683 bc
	SolAxe	M.9	935	23342	2430	223	55,588 bc
	Vertical Axis	B.9	854	20971	2675	227	60,479 bc
	Vertical Axis	G.16	783	19180	2367	218	52,018 c
	Vertical Axis	M.9	1007	24336	2871	224	63,733 bc
	Tall Spindle	B.9	1261	31240	3725	223	84,347 a
	Tall Spindle	G.16	1198	29554	3056	218	66,925 bc
	Tall Spindle	M.9	1243	29381	3254	211	69,101 b
LSD	P≤0.05		279	5965	631	9	15163
McIntosh	Central Leader	MM.111	475	3167	804	164	5,308 g
	Slender Pyramid	G.30	695	3202	2237	148	10,207 f
	Slender Pyramid	M.26	603	3053	1575	150	7,170 g
	SolAxe	B.9	1128	4708	2934	146	13,353 de
	SolAxe	G.16	719	3213	2574	142	10,798 ef
	SolAxe	M.9	1144	4683	3619	149	17,100 bc
	Vertical Axis	B.9	794	2698	2644	143	10,580 ef
	Vertical Axis	G.16	789	3451	2647	146	11,557 ef
	Vertical Axis	M.9	1015	3875	3819	144	14,970 cd
	Tall Spindle	B.9	1602	5800	4792	142	19,396 ab
	Tall Spindle	G.16	1073	6195	3763	143	17,651 bc
	Tall Spindle	M.9	1619	6616	5460	143	21,494 a
LSD	P≤0.05		279	5965	631	9	2964

*Means followed by the same letter do not differ significantly.